


power function  $y = 2x^5$ , and the denominator can be approximated by the power function  $y = x^3$ . This means that as  $x \rightarrow -\infty$  or as  $x \rightarrow \infty$ ,

$$G(x) = \frac{2x^5 - x^3 + 2}{x^3 - 1} \approx \frac{2x^5}{x^3} = 2x^{5-3} = 2x^2$$

Since this is not linear, the graph of  $G$  has no horizontal or oblique asymptote. The graph of  $G$  will behave like  $y = 2x^2$  as  $x \rightarrow \pm\infty$ .

 **Now Work** PROBLEMS 43, 45 AND 47

### 3.4 Assess Your Understanding

'Are You Prepared?'' Answers are given at the end of these exercises. If you get a wrong answer, read the pages listed in red.

- True or False** The quotient of two polynomial expressions is a rational expression. (p. A45)
- What is the quotient and remainder when  $3x^4 - x^2$  is divided by  $x^3 - x^2 + 1$ . (pp. A28–A29)
- Graph  $y = \frac{1}{x}$ . (pp. 15–16)
- Graph  $y = 2(x + 1)^2 - 3$  using transformations. (pp. 89–93)

### Concepts and Vocabulary

- True or False** The domain of every rational function is the set of all real numbers.
- If, as  $x \rightarrow -\infty$  or as  $x \rightarrow \infty$ , the values of  $R(x)$  approach some fixed number  $L$ , then the line  $y = L$  is a \_\_\_\_\_ of the graph of  $R$ .
- If, as  $x$  approaches some number  $c$ , the values of  $|R(x)| \rightarrow \infty$ , then the line  $x = c$  is a \_\_\_\_\_ of the graph of  $R$ .
- For a rational function  $R$ , if the degree of the numerator is less than the degree of the denominator, then  $R$  is \_\_\_\_\_.
- True or False** The graph of a rational function may intersect a horizontal asymptote.
- True or False** The graph of a rational function may intersect a vertical asymptote.
- If a rational function is proper, then \_\_\_\_\_ is a horizontal asymptote.
- True or False** If the degree of the numerator of a rational function equals the degree of the denominator, then the ratio of the leading coefficients gives rise to the horizontal asymptote.

### Skill Building

In Problems 13–24, find the domain of each rational function.

$$13. R(x) = \frac{4x}{x-3}$$

$$14. R(x) = \frac{5x^2}{3+x}$$

$$15. H(x) = \frac{-4x^2}{(x-2)(x+4)}$$

$$16. G(x) = \frac{6}{(x+3)(4-x)}$$

$$17. F(x) = \frac{3x(x-1)}{2x^2-5x-3}$$

$$18. Q(x) = \frac{-x(1-x)}{3x^2+5x-2}$$

$$19. R(x) = \frac{x}{x^3-8}$$

$$20. R(x) = \frac{x}{x^4-1}$$

$$21. H(x) = \frac{3x^2+x}{x^2+4}$$

$$22. G(x) = \frac{x-3}{x^4+1}$$

$$23. R(x) = \frac{3(x^2-x-6)}{4(x^2-9)}$$

$$24. F(x) = \frac{-2(x^2-4)}{3(x^2+4x+4)}$$


In Problems 25–30, use the graph shown to find:

- (a) The domain and range of each function  
(d) Vertical asymptotes, if any


- (b) The intercepts, if any  
(e) Oblique asymptotes, if any

- (c) Horizontal asymptotes, if any


25.




26.




27.




28.



29.



30.



In Problems 31–42, graph each rational function using transformations.

31.  $F(x) = 2 + \frac{1}{x}$

32.  $Q(x) = 3 + \frac{1}{x^2}$

33.  $R(x) = \frac{1}{(x-1)^2}$

34.  $R(x) = \frac{3}{x}$

35.  $H(x) = \frac{-2}{x+1}$

36.  $G(x) = \frac{2}{(x+2)^2}$

37.  $R(x) = \frac{-1}{x^2 + 4x + 4}$

38.  $R(x) = \frac{1}{x-1} + 1$

39.  $G(x) = 1 + \frac{2}{(x-3)^2}$

40.  $F(x) = 2 - \frac{1}{x+1}$

41.  $R(x) = \frac{x^2 - 4}{x^2}$

42.  $R(x) = \frac{x-4}{x}$

In Problems 43–54, find the vertical, horizontal, and oblique asymptotes, if any, of each rational function.

43.  $R(x) = \frac{3x}{x+4}$

44.  $R(x) = \frac{3x+5}{x-6}$

45.  $H(x) = \frac{x^3 - 8}{x^2 - 5x + 6}$

46.  $G(x) = \frac{x^3 + 1}{x^2 - 5x - 14}$

47.  $T(x) = \frac{x^3}{x^4 - 1}$

48.  $P(x) = \frac{4x^2}{x^3 - 1}$

49.  $Q(x) = \frac{2x^2 - 5x - 12}{3x^2 - 11x - 4}$

50.  $F(x) = \frac{x^2 + 6x + 5}{2x^2 + 7x + 5}$

51.  $R(x) = \frac{6x^2 + 7x - 5}{3x + 5}$

52.  $R(x) = \frac{8x^2 + 26x - 7}{4x - 1}$

53.  $G(x) = \frac{x^4 - 1}{x^2 - x}$

54.  $F(x) = \frac{x^4 - 16}{x^2 - 2x}$

### Mixed Practice

In Problems 55–60, for each rational function:

- (a) Graph the rational function using transformations. (b) State the domain and the range.  
(c) State the vertical asymptote and the horizontal asymptote.

55.  $R(x) = \frac{1}{x+3} - 2$

56.  $R(x) = \frac{1}{x-1} + 5$

57.  $R(x) = \frac{-2}{(x-1)^2} + 3$

58.  $R(x) = \frac{3}{(x+2)^2} - 1$

59.  $R(x) = 1 + \frac{4}{x^2 - 2x + 1}$

60.  $R(x) = 2 - \frac{2}{x^2 + 6x + 9}$

## Applications and Extensions

61. **Gravity** In physics, it is established that the acceleration due to gravity,  $g$  (in meters/sec<sup>2</sup>), at a height  $h$  meters above sea level is given by

$$g(h) = \frac{3.99 \times 10^{14}}{(6.374 \times 10^6 + h)^2}$$

where  $6.374 \times 10^6$  is the radius of Earth in meters.

- What is the acceleration due to gravity at sea level?
  - The Willis Tower in Chicago, Illinois, is 443 meters tall. What is the acceleration due to gravity at the top of the Willis Tower?
  - The peak of Mount Everest is 8848 meters above sea level. What is the acceleration due to gravity on the peak of Mount Everest?
  - Find the horizontal asymptote of  $g(h)$ .
  - Solve  $g(h) = 0$ . How do you interpret your answer?
62. **Population Model** A rare species of insect was discovered in the Amazon Rain Forest. Environmentalists protect the species by declaring the insect endangered and transplant the insect into a protected area. The population  $P$  of the insect  $t$  months after being transplanted is

$$P(t) = \frac{50(1 + 0.5t)}{(2 + 0.01t)}$$

- How many insects were discovered? In other words, what was the population when  $t = 0$ ?
  - What will the population be after 5 years?
  - Determine the horizontal asymptote of  $P(t)$ . What is the largest population that the protected area can sustain?
63. **Resistance in Parallel Circuits** From Ohm's law for circuits, it follows that the total resistance  $R_{\text{tot}}$  of two components hooked in parallel is given by the equation

$$R_{\text{tot}} = \frac{R_1 R_2}{R_1 + R_2}$$

where  $R_1$  and  $R_2$  are the individual resistances.

- Let  $R_1 = 10$  ohms, and graph  $R_{\text{tot}}$  as a function of  $R_2$ .
- Find and interpret any asymptotes of the graph obtained in part (a).
- If  $R_2 = 2\sqrt{R_1}$ , what value of  $R_1$  will yield an  $R_{\text{tot}}$  of 17 ohms?

Source: [en.wikipedia.org/wiki/Series\\_and\\_parallel\\_circuits](http://en.wikipedia.org/wiki/Series_and_parallel_circuits)

64. **Newton's Method** In calculus, you will learn that if

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0$$

is a polynomial function, then the *derivative* of  $p(x)$  is

$$p'(x) = n a_n x^{n-1} + (n-1) a_{n-1} x^{n-2} + \cdots + 2 a_2 x + a_1$$

Newton's Method is an efficient method for finding the  $x$ -intercepts (or real zeros) of a function, such as  $p(x)$ . The following steps outline Newton's Method.

**STEP 1:** Select an initial value  $x_0$  that is somewhat close to the  $x$ -intercept being sought.

**STEP 2:** Find values for  $x$  using the relation

$$x_{n+1} = x_n - \frac{p(x_n)}{p'(x_n)} \quad n = 0, 1, 2, \dots$$

until you get two consecutive values  $x_n$  and  $x_{n+1}$  that agree to whatever decimal place accuracy you desire.

**STEP 3:** The approximate zero will be  $x_{n+1}$ .

Consider the polynomial  $p(x) = x^3 - 7x - 40$ .

- Evaluate  $p(5)$  and  $p(-3)$ .
- What might we conclude about a zero of  $p$ ? Explain.
- Use Newton's Method to approximate an  $x$ -intercept,  $-3 < r < 5$ , of  $p(x)$  to four decimal places.
- Use a graphing utility to graph  $p(x)$  and verify your answer in part (c).
- Using a graphing utility, evaluate  $p(r)$  to verify your result.

65. **Exploration** The standard form of the rational function  $R(x) = \frac{mx + b}{cx + d}$  where  $c \neq 0$ , is  $R(x) = a \cdot \left( \frac{1}{x - h} \right) + k$ . To write a rational function in standard form requires long division.

- Write the rational function  $R(x) = \frac{2x + 3}{x - 1}$  in standard form by writing  $R$  in the form

$$\text{Quotient} + \frac{\text{remainder}}{\text{divisor}}$$

- Graph  $R$  using transformations.
  - Determine the vertical asymptote and the horizontal asymptote of  $R$ .
66. **Exploration** See problem 65.

- Write the rational function  $R(x) = \frac{-6x + 16}{2x - 7}$  in the

$$\text{form Quotient} + \frac{\text{remainder}}{\text{divisor}}$$

- Factor out the coefficient on  $x$  in the divisor to write  $R$  in the form  $R(x) = \frac{\text{remainder}}{m(x - h)} + k$ .
- Write the function found in part (b) in the standard form  $R(x) = a \cdot \left( \frac{1}{x - h} \right) + k$ .
- Graph the function found in part (c) using transformations.
- Determine the vertical asymptote and the horizontal asymptote of  $R$ .

## Discussion and Writing

- If the graph of a rational function  $R$  has the vertical asymptote  $x = 4$ , the factor  $x - 4$  must be present in the denominator of  $R$ . Explain why.
- If the graph of a rational function  $R$  has the horizontal asymptote  $y = 2$ , the degree of the numerator of  $R$  equals the degree of the denominator of  $R$ . Explain why.

- Can the graph of a rational function have both a horizontal and an oblique asymptote? Explain.
- Make up a rational function that has  $y = 2x + 1$  as an oblique asymptote. Explain the methodology that you used.

